

Exercises: Stefan Liebler (stefan.liebler@kit.edu) (Office 12/03 - Build. 30.23)

**Exercise 1: Vacuum of the Gupta-Bleuler photon**

**2+4 = 6 points**

In the Gupta-Bleuler formalism of the free photon field the most general vacuum state reads

$$|\phi\rangle = \sum_{n=0}^{\infty} C_n |\phi_n\rangle.$$

The states  $|\phi_n\rangle$  do not include transverse photons, but exactly  $n$  scalar and longitudinal photons. The additional condition

$$(a_3(k) - a_0(k))|\phi_n\rangle = 0$$

makes them physical states. We moreover choose  $|\phi_0\rangle = |0\rangle$ .

- (a) Show that the most general form of  $|\phi_1\rangle$  is given by

$$|\phi_1\rangle = \int d\tilde{q} f(q) (a_3^\dagger(q) - a_0^\dagger(q)) |0\rangle.$$

*Hint:* Make the ansatz  $|\phi_1\rangle = \int d\tilde{q} \sum_{r=0,3} a_r^\dagger(q) f_r(q) |0\rangle$ .

- (b) Show that the expectation value of the photon field in the above general vacuum state corresponds to a gauge fixing, i.e.

$$\langle\phi|A_\mu(x)|\phi\rangle = \partial_\mu\Lambda(x),$$

where the function  $\Lambda(x)$  using the explicit polarisation vectors  $\varepsilon_0^\mu(k) = n^\mu$  and  $\varepsilon_3^\mu(k) = \frac{k^\mu}{k \cdot n} - n^\mu$  is given by

$$\Lambda(x) = \int \frac{d\tilde{k}}{k \cdot n} 2\text{Re} (iC_0^* C_1 e^{-ik \cdot x} f(k)).$$

Therein  $f(k)$  is identical to the one in subexercise (a). The function  $\Lambda(x)$  fulfills  $\square\Lambda(x) = 0$  and can be chosen arbitrarily through the choice of the corresponding vacuum state  $|\phi\rangle$ .

*Hint:* First show that  $\langle\phi_n|N A_\mu(x)|\phi_{n-1}\rangle = \langle\phi_n|A_\mu(x)|\phi_{n-1}\rangle$  with

$$N = \int d\tilde{k} (a_3^\dagger(k)a_3(k) - a_0^\dagger(k)a_0(k))$$

counting longitudinal and scalar photons. Thus it yields  $\langle\phi_n|A_\mu(x)|\phi_{n-1}\rangle = 0$  for  $n \neq 1$ .

**Exercise 2: Massive vector boson****2+3+4 = 9 points**

We consider a vector boson with mass  $m \neq 0$ , which enters the Lagrangian density as follows

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{m^2}{2}A_\mu A^\mu$$

with  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ .

- (a) Derive the equation of motion for  $A^\mu$ , that is known as Proca equation. Show that this equation necessarily implies  $\partial_\mu A^\mu = 0$  and that, once this condition is imposed, it is equivalent to a set of massive Klein-Gordon equations, namely one for each of the non-vanishing components of  $A^\mu$ .
- (b) We introduce a Fourier decomposition for the massive vector boson in analogy to the photon field, i.e.

$$A^\mu(x) = \int d\vec{k} \sum_{r=0}^3 (\varepsilon_r^\mu(k) a_r(k) e^{-ik \cdot x} + \varepsilon_r^{\mu*}(k) a_r^\dagger(k) e^{ik \cdot x}) .$$

A priori, this includes four polarization vectors  $\varepsilon_r^\mu(k)$ . Due to  $\partial_\mu A^\mu = 0$  and thus  $\sum_r k_\mu \varepsilon_r^\mu(k) a_r(k) = 0$  only three polarization vectors are physical. Show that a convenient basis for these polarization vectors, in the reference frame with  $\vec{k} = (0, 0, |\vec{k}|)$ , is given by

$$\varepsilon_1 = (0, 1, 0, 0), \quad \varepsilon_2 = (0, 0, 1, 0), \quad \varepsilon_3 = \frac{1}{m}(|\vec{k}|, 0, 0, \omega_k)$$

for the three physical polarization vectors, which are orthogonal to the unphysical polarisation vector  $\varepsilon_0^\mu = k^\mu/m$ . The physical polarization vectors obey the orthonormality condition  $\varepsilon_r^\mu(k) \varepsilon_{\mu s}^*(k) = -\delta_{rs}$ . Using these explicit expressions show that the completeness relation of the physical polarization vectors reads

$$\sum_{r=1}^3 \varepsilon_r^\mu(k) \varepsilon_r^{\nu*}(k) = -g^{\mu\nu} + \frac{k^\mu k^\nu}{m^2} .$$

*Hint:* In the rest frame of the particle,  $k' = (m, 0, 0, 0)$ , we e.g. choose  $\varepsilon'_0 = (1, \vec{0})$  and the three unit vectors  $\varepsilon'_i = (0, \vec{e}_i)$ . Then the vectors  $\varepsilon'_i$  automatically fulfill  $k'_\mu \varepsilon_i^{\prime\mu} = 0$ . Boost from the rest frame into the above reference frame. *Add-on:* We showed the completeness relation in a special frame, but it is actually Lorentz-covariant.

- (c) We now impose standard bosonic commutation relations for the surviving operators. They read

$$\begin{aligned} [a_r(k), a_s(k')] &= [a_r^\dagger(k), a_s^\dagger(k')] = 0, \\ [a_r(k), a_s^\dagger(k')] &= \delta_{rs} 2\omega_k (2\pi)^3 \delta^{(3)}(\vec{k} - \vec{k}') . \end{aligned}$$

Verify that the propagator of the massive vector boson takes the form

$$\langle 0|T A^\mu(x) A^\nu(y)|0\rangle = \int \frac{d^4k}{(2\pi)^4} \frac{i}{k^2 - m^2 + i\epsilon} \left( -g^{\mu\nu} + \frac{k^\mu k^\nu}{m^2} \right) e^{-ik(x-y)} .$$

*Add-on:* As the photon has only two rather than three physical degrees of freedom, the limit  $m \rightarrow 0$  of this propagator is not well-defined and does not yield the photon propagator.