

Einführung in Theoretische Teilchenphysik

Lecture: PD. Dr. S Gieseke – Exercises: Dr. D. López-Val, Dr. S. Patel, Dr. M. Rauch

Exercise Sheet 1

Submission: Mo, 23.10.17, 12:00

Discussion: Mon, 23.10.17 14:00 Room 11/12
Wed, 25.10.17 09:45 Room 10/1

Exercise 1: Warm up: Natural Units

The so-called *Natural Units* ($\hbar = c = 1$) are commonly employed in Particle Physics. This system of units implies that velocities are measured in units of speed of light c ; while forces are measured in units of Planck's constant (divided by 2π) \hbar .

- (a) How do Meters (length), Seconds (time), and MeV (energies) relate to each other in this system?

Hint: $c \approx 2,99 \times 10^8$ m/s, und $\hbar = 6,58 \times 10^{-22}$ MeV s.

- (b) What is the mass (in Kilogramm) corresponding to 1 MeV?

Hint: $1\text{eV} = 1,60 \times 10^{-19}$ J.

- (c) Muons μ^\pm have a mass of $m_{\mu^\pm} = 1,88 \times 10^{-28}$ kg and a lifetime of $\tau_\mu = 2,2 \times 10^{-6}$ s. In turn, the mass of the W bosons is $m_W = 1,43 \times 10^{-25}$ kg and their corresponding decay rate yields $\tau_W = 3,1 \times 10^{-25}$ s. Express (in MeV) both the mass and the total decay width $\Gamma = 1/\tau$ for each particle, and compute the respective width/mass ratios Γ/m . Which of the two particles can you conclude to be more stable?

Exercise 2: Energy and distance scales

- (a) Estimate the typical length scale that is probed by experiments:
(i) at the *Large Electron-Positron Collider* (LEP) ($\sqrt{S} = 45 - 209$ GeV);
(ii) at the Run II Large Hadron Collider ($\sqrt{S} = 13$ TeV)

In view of these results: why are the notions of *Particle Physics* and *High Energy Physics* so often used as synonyms?

Hint: Recall the notion of *wave-particle duality* and how it is implemented by the *de Broglie relation*

- (b) Estimate the energy necessary to test the characteristic length scales of a quantum theory of gravitation.

Hint: Use *dimensional analysis*, and bear in mind what are the key scales characterizing *Gravity* ($G_N = 667408(31) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$); a *quantum regime* ($\hbar = 6.62607004 \times 10^{-34} \text{ m}^2 \text{ kg s}^{-1}$); and *relativistic conditions* ($c = 299792458 \text{ m s}^{-1}$).

Exercise 3: Range of the fundamental interactions

According to the current paradigm of Particle Physics, the interactions between two particles are described by the exchange of bosonic *force carriers*. These particles can be massive, and in general propagate off-shell ($p_\mu p^\mu \neq m^2$).

- (a) The effective (residual) Strong Interaction between two nucleons is mediated by the interchange of virtual (off-shell) *Pions* (these are quark-antiquark bound states), and its typical range is of the order of 1 fm. Recalling the Heisenberg uncertainty relations, and assuming that the intermediate pions propagate at the speed of light, estimate the pion mass and compare it to its actual value.
- (b) Using similar arguments, estimate the typical range of:
- (i) The weak interaction (the masses of the W and Z bosons are respectively $M_W \approx 80 \frac{\text{GeV}}{c^2}$ and $M_Z \approx 91 \frac{\text{GeV}}{c^2}$).
 - (ii) The Electromagnetic interaction;
 - (iii) The Strong interaction. Why is your result somehow counterintuitive in the latter case?