

Exercises: Stefan Liebler (stefan.liebler@kit.edu) (Office 12/03 - Build. 30.23)  
Martin Gabelmann (martin.gabelmann@kit.edu) (Office 12/17 - Build. 30.23)  
Jonas Müller (jonas.mueller@kit.edu) (Office 12/17 - Build. 30.23)

### Exercise 1: Propagator of a free particle

We want to evaluate the path integral of a free particle in one-dimension by parametrizing all potential paths through a variation of the coefficients of a Fourier series. The propagator is defined by

$$\langle q_f, t_f | q_i, t_i \rangle = \int \mathcal{D}q e^{\frac{i}{\hbar} S} \quad \text{with} \quad S = \int_{t_i}^{t_f} dt \frac{m}{2} \dot{q}^2$$

with the boundary conditions  $q(t_i) = q_i$  and  $q(t_f) = q_f$ .

- Write down the classical, linear path of the particle  $q_c(t)$  as a function of  $q_i, q_f, t_i$  and  $t_f$ .
- In order to evaluate the path integral we allow for quantum mechanical perturbations around the classical path,  $q(t) = q_c(t) + \delta q(t)$ , and expand  $\delta q(t)$  in a Fourier series knowing that the fluctuations vanish at  $q_i$  and  $q_f$ . We therefore obtain

$$\delta q(t) = \sum_{n=1}^{\infty} a_n \sin \left[ \frac{n\pi}{t_f - t_i} (t - t_i) \right].$$

Using the orthogonality of different modes, show that the action  $S$  turns into

$$S = \frac{m}{2} \frac{(q_f - q_i)^2}{t_f - t_i} + \frac{m}{2} \sum_{n=1}^{\infty} \frac{1}{2} \frac{(n\pi)^2}{t_f - t_i} a_n^2.$$

- In this example the path integral can be rewritten in terms of integrals over the spectrum of the Fourier coefficients, such that  $\int \mathcal{D}q = c \int \prod_{n=1}^{\infty} da_n$  with a normalization constant  $c$ . Rearrange the integrals and show

$$\langle q_f, t_f | q_i, t_i \rangle = \tilde{c}(t_f - t_i) \exp \left[ \frac{i}{\hbar} \frac{m}{2} \frac{(q_f - q_i)^2}{t_f - t_i} \right],$$

where the normalization constant  $\tilde{c}(t_f - t_i)$  can only be a function of the time difference.

*Hint:* You can use  $\int_{-\infty}^{\infty} da \exp \left[ \frac{i}{\hbar} \frac{m}{2} \frac{(n\pi)^2}{t_f - t_i} a^2 \right] = \left( -\frac{imn^2\pi}{2\hbar(t_f - t_i)} \right)^{-1/2}$ .

- Show that the normalization constant  $\tilde{c}(t)$  is given by

$$\tilde{c}(t) = \sqrt{\frac{m}{2\pi i \hbar t}} \quad \text{from the requirement} \quad \int dq \langle q_f, t_f | q, t \rangle \langle q, t | q_i, t_i \rangle = \langle q_f, t_f | q_i, t_i \rangle.$$

*Hint:* You can use  $\int_{-\infty}^{\infty} dq \exp \left[ \frac{i}{\hbar} \frac{m}{2} \left( \frac{(q_f - q)^2}{(t_f - t)} + \frac{(q - q_i)^2}{(t - t_i)} \right) \right] = \sqrt{\frac{2\pi i \hbar (t_f - t)(t - t_i)}{m(t_f - t_i)}} \exp \left[ \frac{i}{\hbar} \frac{m}{2} \frac{(q_f - q_i)^2}{t_f - t_i} \right]$ .

- The wave-function of the particle thus evolves according to

$$\psi(q, t) = \int dq_i \langle q, t | q_i, 0 \rangle \psi(q_i, 0).$$

Show that  $\psi(q, t)$  satisfies the Schrödinger equation and that  $\psi(q, t) \rightarrow \psi(q, 0)$  for  $t \rightarrow 0$ .

*Hint:* Fresnel representation of the  $\delta$  distribution.

## Exercise 2: Path integral formalism for the harmonic oscillator

We investigate the propagator  $\langle q_f, t_f | q_i, t_i \rangle$  of the one-dimensional harmonic oscillator in the path integral formalism of quantum mechanics. In contrast to the previous exercise we use  $\hbar = 1$ .

- (a) Show that due to the completeness relation of the energy eigenstates  $|n\rangle$  with energy  $E_n$  we can write

$$\langle q_f, T | q_i, 0 \rangle = \sum_n \Phi_n(q_f) \Phi_n^*(q_i) e^{-iE_n T} \quad (1)$$

with  $\Phi_n(q) = \langle q | n \rangle$ .

- (b) The propagator of the harmonic oscillator can be determined from the path integral formalism and yields

$$\langle q_f, T | q_i, 0 \rangle = \sqrt{\frac{m\omega}{2\pi i \sin \omega T}} \exp \left\{ \frac{im\omega}{2 \sin \omega T} [(q_f^2 + q_i^2) \cos \omega T - 2q_i q_f] \right\}. \quad (2)$$

Determine the wave function of the ground state  $\Phi_0(q)$ . In order to do so, set the time to an imaginary value  $T = -it, t \rightarrow \infty$ , rewrite sin and cos in terms of exponentials and compare Eq. 1 and Eq. 2. For this purpose rewrite Eq. 2 in the form  $e^{-\omega/2t} f(e^{-\omega t})$  and expand around the argument of  $f(x)$ . The first term corresponds to the ground state. *Note:* We refer to the literature for the derivation of the path integral representation of the harmonic oscillator.

- (c) Calculate the wave function of the first excited state  $\Phi_1(q)$ .
- (d) Show that the energy levels are given by  $E_n = \omega(n + \frac{1}{2})$ .  
*Hint:* Again compare Eq. 1 and Eq. 2.
- (e) We finally allow the harmonic oscillator to be influenced by an external driving force  $J(t)$ . The propagator then takes the form

$$\langle q_f, t_f | q_i, t_i \rangle_J = \int \mathcal{D}q \exp \left[ i \int_{t_i}^{t_f} dt \left( \frac{1}{2} \dot{q}^2 - \frac{1}{2} \omega^2 q^2 + J(t)q(t) \right) \right] \quad (3)$$

with the boundary conditions  $q(t_i) = q_i$  and  $q(t_f) = q_f$ .

- (f) Show that for  $\omega^2 \rightarrow \omega^2 - i\epsilon$  the propagator in Eq. 3 can be rewritten in the form

$$Z[J] = \langle q_f, \infty | q_i, -\infty \rangle_J = \langle q_f, \infty | q_i, -\infty \rangle_0 \exp \left[ -\frac{i}{2} \int dE \frac{\tilde{J}(E) \tilde{J}(-E)}{E^2 - \omega^2 + i\epsilon} \right], \quad (4)$$

where we have introduced

$$\tilde{J}(E) = \int_{-\infty}^{\infty} \frac{dt}{\sqrt{2\pi}} e^{-iEt} J(t) \quad \leftrightarrow \quad J(t) = \int_{-\infty}^{\infty} \frac{dE}{\sqrt{2\pi}} e^{iEt} \tilde{J}(E).$$

*Hints:* Make use of a Fourier transform ( $t \leftrightarrow E$ ) in the exponent of Eq. 3 for both  $J(t)$  and  $q(t)$ , write  $Jq = \frac{1}{2}[Jq + qJ]$ , use  $\int dt \exp[i(E + E')t] = 2\pi\delta(E + E')$  and motivate a shift  $\tilde{q}(E) \rightarrow \tilde{q}(E) - \frac{\tilde{J}(E)}{E^2 - \omega^2}$ .

- (g) Transform the exponent in Eq. 4 into a time integral, i.e.

$$-\frac{i}{2} \int dt dt' J(t) \Delta(t - t') J(t') \quad \text{with} \quad \Delta(t - t') = \int \frac{dE}{2\pi} \frac{e^{-iE(t-t')}}{E^2 - \omega^2 + i\epsilon},$$

and discuss its physical meaning.