KA-TP-2-'97
May '97

On Sums of Powers of Zeros of Polynomials *

Wolfdieter L a n g **

Institut für Theoretische Physik
Universität Karlsruhe
Kaiserstrasse 12, D-76128 Karlsruhe
Germany

ABSTRACT

Due to Girard's (sometimes called Waring's) formula the sum of the $r-$th power of the zeros of every one variable polynomial of degree N, $P_N(x)$, can be given explicitly in terms of the coefficients of the monic $\tilde{P}_N(x)$ polynomial. This formula is closely related to a known $N-1$ variable generalisation of Chebyshev's polynomials of the first kind, $T_r^{(N-1)}$. The generating function of these power sums (or moments) is known to involve the logarithmic derivative of the considered polynomial. This entails a simple formula for the Stieltjes transform of the distribution of zeros. Perron-Stieltjes inversion can be used to find this distribution, *e.g.* for $N \to \infty$.

Classical orthogonal polynomials are taken as examples. The results for ordinary Chebyshev $T_N(x)$ and $U_N(x)$ polynomials are presented in detail. This will correct a statement about power sums of zeros of Chebyshev's $T-$polynomials found in the literature. For the various cases (Jacobi, Laguerre, Hermite) these moment generating functions provide solutions to certain Riccati equations.

AMS MSC numbers: 30C15, 33C50, 34L30
PACS numbers: 02.50.Sk, 02.10.Nj 02.30.Gp

* The complete ps-file is also available via www under
http://www-itp.physik.uni-karlsruhe.de/~wl/preprints.html

** E-mail address: wolfdieter.lang@physik.uni-karlsruhe.de